

# New Trends on Algebra of Operators and Tensor Products

U. K. Srivastava<sup>1</sup>, L. R. Pratap<sup>2</sup>, A. K. Singh<sup>3</sup>, MD. Shahabuddin<sup>4</sup>

*Department of Mathematics*

*R.S.S. College Chochahan , P.O. :- Aniruddh Belsar, Dist. – Muzaffarpur – 844111, B.R.A. Bihar University, Muzaffarpur – 842001, Bihar, India.*

*M.S.K.B. College, Muzaffarpur, 842001, B.R.A. Bihar University, Muzaffarpur – 842001, Bihar, India.*

*S.N.S. College, Hajipur (Vaishali ), - 844101 , B.R.A. Bihar University, Muzaffarpur – 842001, Bihar, India.*

*P.R.R.D. College, Bairgania, Sitamarhi – 843313, B.R.A. Bihar University, Muzaffarpur – 842001, Bihar, India.*

**Abstract--** This paper presents the study of Algebra of operators and Tensor products . Here, we consider  $\mathbb{R}$  as additive group of reals with discrete topology and several ways of constructing  $C^*$  - algebras Canonically associated with  $\mathbb{R}$  and  $\pi$  , the Universal representation of  $\mathbb{R}$  on Hilbert space  $H$ , it is proved in this paper that all  $C^*$  - algebras homomorphism and representation will be  $*$  - preserving.

**Keywords--** Hilbert space, Tensor product,  $C^*$  - tensor norms,  $C^*$  - algebras , Normal and Binormal norms ,  $W^*$ -algebras.

## I. INTRODUCTION

E.G. EFFROS (1) and Kothe (3,4) are the pioneer workers of the present area . In fact , the present work is the extension of work done by Halub , J.R. (2) , Kumar et al. (5), Kumar et al. (6) , Kumar et al. (7) , Srivastava et al. (8) , Srivastava et al. (9), Srivastava et al. (10) and Srivastava et al. (11). In this paper , we have studied new trends on Algebra of operators and Tensor products.

### *Mathematical Treatment Of The Problem*

Let,  $\mathbb{R}$  = additive group of reals with discrete topology. There are several ways of constructing  $c^*$ -algebras canonically associated with  $\mathbb{R}$ . For example if  $\pi$  is the universal representation of  $\mathbb{R}$  on the Hilbert space  $H$ , the group  $c^*$ - algebra  $c^*(\mathbb{R})$  is the  $c^*$ -subalgebra of  $\mathcal{L}(H)$  generated by the set  $\{\pi(g) : g \in \mathbb{R}\}$ . The left regular group  $c^*$ - algebra  $c_l^*(\mathbb{R})$  is the  $c^*$ -subalgebra of  $\mathcal{L}(\ell^2(\mathbb{R}))$  generated by  $\pi\ell(\mathbb{R})$ ,  $\pi\ell$  being the left regular representation of  $\mathbb{R}$ , when

$$\pi\ell(g)\xi(h) = \xi(g^{-1}(h)), \quad (g, h \in \mathbb{R}, \xi \in \ell^2(\mathbb{R}))$$

The right regular group  $C^*$ -algebra  $C_r^*(\mathbb{R})$  is defined analogously using the right regular representation  $\pi_r$  of  $\mathbb{R}$ , when

$$\pi_r(g)\xi(h) = \xi(hg), \quad (g, h \in \mathbb{R}, \xi \in \ell^2(\mathbb{R}))$$

By its definition  $\pi$  contains  $\pi_l$  and  $\pi_r$  , and there are natural homomorphism  $\lambda_l$  and  $\lambda_r$  of  $c^*(\mathbb{R})$  onto  $c_l(\mathbb{R})$  and  $c_r^*(\mathbb{R})$  resp.

(Note also that the representations  $\pi_l$  and  $\pi_r$  are equivalent, so that  $c_l^*(\mathbb{R})$  and  $c_r^*(\mathbb{R})$  are infact isomorphic).

Let  $A$  and  $B$  be  $c^*$ -algebras, with algebraic tensor product  $A \otimes B$ . In general there are several distinct (usually incomplete).

$c^*$ - norms on  $A \otimes B$ . Two such norms are of particular interest: the maximal norm  $\nu$  of Guichander and the minimal (or spatial) norm  $\alpha$  of Turumaru.

If  $\pi_1$  and  $\pi_2$  are representations of  $A$  and  $B$ , respectively, on the Hilbert space  $H$ ,  $\{\pi_1, \pi_2\}$  is said to be a committing pair of representations of  $A, B$  if

$$\pi_1(a)\pi_2(b) = \pi_2(b)\pi_1(a) \quad (a \in A, b \in B)$$

The norm  $\nu$  is defined by

$$\nu(\sum_i a_i \otimes b_i) = \sup (|| \sum_i \pi_1(a_i) \pi_2(b_i) || )$$

the supremum being taken over all commuting pairs of representations of  $A, B$ . The norm  $\alpha$  is defined as follows:

if  $x \in A \otimes B$ ,  $\alpha(x)$  is the smallest non-negative real number  $\mathcal{R}$  such that

$$\langle f \otimes g, a^*x^*xa \rangle \leq \mathcal{R}^2 \langle f \otimes g, a^*a \rangle$$

For all  $a \in A \otimes B$  and all sates  $f$  and  $g$  of  $A$  and  $B$  respectively. If for all  $\beta\alpha = \nu$  on  $A \otimes B$ ,  $A$  is said to be nuclear (the terminology in due to Lance, whives an introduction to the theory of  $c^*$ - tensor product. For a discrete group  $\mathbb{R}$ ,  $c_l^*(\mathbb{R}) = c^*(\mathbb{R})$  iff  $G$  is amenable, and this is the case iff  $c_l^*(G)$  is nuclear.

Let  $\mathbb{R}$  be a discrete group and let be the representation of  $c^*(\mathbb{R}) \otimes C^*(\mathbb{R})$  on  $\ell^2(\mathbb{R})$  given by

$$\lambda(\sum a_i \otimes b_i) = \sum \lambda_l(a_i) \lambda_r(b_i)$$

It is natural to ask relative to which  $c^*$ - norms  $\eta$  on  $c^*(R) \odot c^*(R)$   $\lambda$  is bounded. If  $\eta = \alpha$ ,  $\lambda$  is already bounded with  $\eta = \alpha$ , the same is true if  $R$  is amenable (by the remark above). One of the main results of this section is essentially that  $R = F_2$ , the free group on two generators,  $\lambda$  is bounded when  $\eta = \alpha$ . A consequence of this fact is that, even in the separable case, it is not always possible to describe the ideal structure of the spatial tensor product of two  $c^*$ -algebras in terms of the ideal structures of the individual algebras. It follows moreover that the spatial  $c^*$ -norm is not in general preserved by quotients, (by quotients respect the maximal norm  $v$ ).

Effros and Lance have recently introduced two new  $c^*$ -tensor norms, the normal and binormal norms. Let  $A$  and  $B$  be  $c^*$ -algebras.

If  $A$  is a  $W^*$ -algebra, the left normal norm  $v_l$  on  $A \odot B$  is defined by

$$v_l(\sum a_i \otimes b_i) = \sup (\| \sum \pi_1(a_i) \pi_2(b_i) \|)$$

The supremum being taken over all commuting pairs of representations  $\{\pi_1, \pi_2\}$  of  $A, B$  with  $\pi_1$  normal. If  $B$  is a  $W^*$ -algebra, the right normal norm  $v_r$  is defined analogously with  $\pi_2$  rather  $\pi_1$  required to be normal. If  $A$  and  $B$  are both  $W^*$ -algebras, the binormal norm  $\beta$  is given by a similar expression, the supremum being taken this time over all commuting pairs of normal representations of  $A, B$ . The notation and terminology used here follow those of [1] for the most part.

If  $A$  and  $B$  are  $c^*$ -algebras and  $\eta$  is a  $c^*$ -norm on  $A \odot B$ , then  $\eta$ -completion of  $A \odot B$  will be denoted  $A \otimes_\eta B$ , the  $\alpha$ -completion will be denoted simply by  $A \otimes B$ . If  $B$  is a  $c^*$ -subalgebra of  $A$ , a linear map  $\rho : A \rightarrow B$  is a retraction if it is a projection of norm 1, i.e. if  $\|\rho\| = 1$  and  $\rho(x) = x$  for  $x \in B$ . Finally, all  $c^*$ -algebras homomorphism and representations will be assumed to be  $*$ -preserving.

1. We recall that a  $W^*$ -algebra  $M$  is said to have the extension property (or to be injective(iii)) if it satisfies the following equivalent condition:

(i) For some faithful normal representation  $\pi$  of  $M$  on the Hilbert space  $H$  there is retraction

$$\rho : \mathcal{L}(H) \rightarrow \pi(M).$$

(ii) such a  $\rho$  exists for every normal representation  $\pi$  of  $M$ .

(iii) For some faithful normal representation  $\pi$  of  $M$  on  $H$  there is a retraction,

$$\rho' : \mathcal{L}(H) \rightarrow \pi(M)'$$

A  $c^*$ -algebra  $A$  is of type E if for every representation  $\pi$  of  $A$ , the weak closure  $\overline{\pi(A)}$  has the extension property (i.e. if  $A^{**}$  is injective).

(2) It is not known whether a  $c^*$ -subalgebra of a  $c^*$ -algebra of type E is automatically of type E also. For certain types of subalgebra this is the case, which is the Main Result as

Let  $A$  be a  $c^*$ -algebra of type E. If  $B$  is a unital  $c^*$ -subalgebra of  $A$  and there is a retraction

$$\rho : A \rightarrow B \text{ then } B \text{ is of type E.}$$

*Proof:* Let  $A^{**}$  non-degenerately as a Von Neumann algebra on the Hilbert space  $H$ . The map  $\rho^{**} : A^{**} \rightarrow B^{**}$  is normal regarding  $B^{**}$  as a  $W^*$ -subalgebra of  $A^{**}$  under its canonical embedding,  $\rho^{**}$  is a retraction. Let  $\sigma$  be a representation of  $B$  on the Hilbert space  $K$ , it is sufficient to assume that  $\sigma$  is non-degenerate. There is a central projection,

$$F \in B^{**} \text{ such that } B^{**} F \cong \overline{\sigma(B)}$$

By hypothesis there is a retraction,

$$\tau : \mathcal{L}(H) \rightarrow A^{**}$$

define  $W : \mathcal{L}(FH) \rightarrow B^{**}$  by,

$$W(T) = (\rho^{**} \odot \tau)(TF) \quad (T \in \mathcal{L}(FH))$$

It is clear that  $\|W\| \leq 1$  and by the nodule property of retractions,

$$W(TF) = W(T)F \text{ for } T \in \mathcal{L}(FH)$$

so that

$$W(T) \in B^{**} F \text{ for } T \in \mathcal{L}(FH)$$

and moreover,

$$W(T) = T \text{ for } T \in B^{**} F.$$

Thus,  $W$  is a retraction with image

$B^{**} F$ . Thus,  $\overline{\sigma(B)}$  has the extension property, so that  $B$  is of type E.

Hence the result.

*Acknowledgment*

The authors are thankful to Prof.(Dr.) S.N. Jha , Ex. Head , Prof. (Dr.) P.K.Sharan, Ex.Head, and Prof.(Dr.) B.P. Kumar, present Head of the Deptt. Of Mathematics, B.R.A.B.U. Muzaffarpur, Bihar , India and Prof. (Dr.)T.N. Singh, Ex. Head ,Ex. Dean (Science) and Ex. Chairman, Research Development Council , B.R.A.B.U. Muzaffarpur, Bihar , India for extending all facilities in the completion of the present research work.

**REFERENCES**

- [1] E.G. EFFROS and E.C. LANCE . (1970) : Tensor Products of Operator Algebras.
- [2] Halub, J.R. (1970 ) : Tensor Product Mapping Math Ann ., Vol. 188 , pp 01-12
- [3] Kothe , G. (1969) : Topological Vector Spaces , Springer – Verlag , I, New York.
- [4] Kothe, G. (1979) : Topological Vector Spaces , Springer – Verlag , II, New York.
- [5] Kumar , N.,Choudhary , D; :International Journal of Emerging Technology Singh , A.K., and Srivastava, and Advanced Engineering Vol. 5(11), pp 177-179. U.K., (2015)
- [6] Kumar Nirmal ; Prasad , C.S; : International Journal of Emerging Technology Shahabudin, Md; and and Advanced Engineering Vol. 6(10), pp 149-152. Srivastava, U.K., (2016)
- [7] Kumar Nirmal ; Choudhary , : International Journal of Emerging Technology D., Talukdar , A.U., and Advanced Engineering Vol. 6(11), pp 170-173. Srivastava, U.K., (2016)
- [8] Srivastava , U.K.,Kumar, S. , : J.P.A.S (Mathematical Science) ,Vol. 17, and Singh , T.N. (2011) pp 188-191.
- [9] Srivastava , U.K., : J.P.A.S (Mathematical Science) ,Vol. 18, Kumar, N., Kumar ,S., pp 189- 191. and Singh, T.N. (2012)
- [10] Srivastava , U.K., kumar , : Proc. Of Math. Soc., B.H.U. Vol. 28, pp 25-28. N., Kumar , S., and Singh, T.N. (2012)
- [11] Srivastava, U.K., Talukdar, : International Journal of Emerging Technology A.U., Shahabuddin, Md., and Advanced Engineering Vol. 6(6), pp 247-250. and Pandit, A.S., (2016)