

Effects of Dietary Lysine on Growth Performance, Carcass Composition and Immunological Responses to Influenza Vaccination in Ac Chickens

Lam Thai Hung¹, Nguyen Trong Ngu²

¹PhD, Agriculture and Aquaculture School, Tra Vinh University, Tra Vinh Province, Vietnam

²PhD, College of Agriculture and Applied Biology, Can Tho University, Can Tho City, Vietnam

Abstract— The aim of this study was to evaluate the effects of different lysine levels on growth, carcass composition and immunological response to bird flu vaccination in Ac chicken. Four hundred fifty one-day-old chicks and two hundred forty of twenty eight-day-old chicken were arranged in a completely randomized design with 5 diets (different lysine contents of 0, 5, 10, 15 and 20% of the diet) and 3 replicates. Birds were fed *ad libitum* and had free access to water. Blood samples were collected to analyze immunological responses to H5N1 vaccination at 28 and 56 day-old. Six birds per a treatment were slaughtered to evaluate bodyweight, carcass composition and chemical composition. Results showed that feed consumption of birds was similar (99.2-111.9 g/bird/week) when increasing dietary lysine, but bodyweight gain and feed conversion ratio were improved ($P<0.05$) in diet with 10% lysine inclusion (52.4 g/bird/week and 2.09 kg feed/kg gain, respectively). In addition, bodyweight, carcass composition and amino acid content of breast muscle of Ac chicken were not different ($P<0.05$) between treatments, but ratio of weight of organs producing immunological antibodies of chicken was greater in the high lysine diets.

Keywords— antibody, growth, local chicken, lysine, performance

I. INTRODUCTION

Lysine is one of the limiting amino acids in broiler diets and it is used to calculate the remaining essential amino acids in the ration [3]. Lysine requirement of broilers is higher in low protein diets for maximum weight gain and feed efficiency [10]. As widely described, increasing dietary lysine generally results in improved feed intake, feed conversion ratio and body weight gain [22]. Moreover, lysine has impact not only on birds' growth but also on their ability against bird flu (H5N1) infection. Previously, [15] found that addition of lysine at high levels to the diets can stimulate insulin secretion from pancreas by aggregating in plasma, which in turn releases amino acid and fatty acids from the bodily saved sources and leads to protein synthesis.

Supportably, the report of [21] indicated that the addition and improvement of lysine in poultry diets improved birds' immune ability against different diseases.

In Vietnam, Ac chicken is an indigenous breed with black skin, meat and bones and is widely adopted in the Mekong Delta. Its meat quality is considered juicy with high levels of amino acids. Similar to other breeds, Ac chickens need amino acids for body growth and protein synthesis, of which lysine is the most important amino acid to concern. Previous work has focused on conservation [26], production [18] and protein and energy levels on performance and egg quality [17] but little was known on the effects of lysine inclusion levels on Ac chicken. This study was therefore designed to determine growth performance, carcass composition and immunological response to influenza vaccination in Ac chickens supplemented with lysine at different content in the diet.

II. MATERIALS AND METHODS

A. Experimental design and diets

The feeding experiment was allocated in a completely randomized design with 5 treatments and 3 replicates (thirty and sixteen birds for each treatment at 0-4 and 5-8 weeks of age, respectively). Vaccination against Newcastle disease was performed at the 5th day of age by nostril and eye drop method and bird flu at the 14th day of age by breast intramuscular injection. The diets were offered *ad libitum* and water was freely available throughout eight weeks. The experimental birds were fed with five diets including control (dietary lysine 1% and 0.9% of 0-4 and 5-8 week-old period, respectively) and other four diets were added with 5%, 10%, 15%, and 20% lysine as compared with the control diet. Details of nutritional values and composition of diets of two phases are shown in Table 1, Table 2 and Table 3, respectively.

TABLE 1
Nutrient value of yellow corn and soybean meal

Feedstuff	Nutrient value (% of feed)						
	DM	ME, kcal/kg	CP	Lys	Met+cys	Ca	P
Yellow corn	91.54	3,222	9.17	0.247	0.779	0.63	0.29
Soybean meal	90.96	2,500	44.96	1.289	1.967	0.63	0.67

DM: dry matter; ME: metabolizable energy; CP: crude protein;
Lys: lysine; Met+cys: methionine + cysteine.

TABLE 2
Experimental diets of birds in 0-4 week of age

Feedstuffs	Treatments				
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20
Yellow corn, g	62.95	63.02	63.07	63.07	63.13
Soybean meal, g	32.93	32.82	32.7	32.63	32.5
Soybean oil, g	0.55	0.55	0.56	0.58	0.6
HCl-Lysine, g	0.43	0.47	0.53	0.58	0.63
DCP, g	1.95	1.95	1.95	1.95	1.95
Shell, g	0.46	0.46	0.46	0.46	0.46
Premix, g	0.25	0.25	0.25	0.25	0.25
Salt, g	0.48	0.48	0.48	0.48	0.48
Nutrient value					
ME, kcal/kg	2,900	2,900	2,900	2,900	2,900
CP, %	21	21	21	21	21
Lysine, %	1	1.05	1.1	1.15	1.2
Ca, %	1.2	1.2	1.2	1.2	1.2
P, %	0.78	0.78	0.78	0.78	0.78

Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding lysine 5%, 10%, 15%, and 20% compared with control treatment; DCP: Dicarboxate phosphorus; CP: crude protein; ca: Calcium; P: phosphorus.

TABLE 3
Experimental diets of birds in 5-8 week of age

Feedstuffs	Treatments				
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20
Yellow corn, g	72.0	72.0	72.0	72.0	72.0
Soybean meal, g	24.46	24.41	24.37	24.33	24.28
HCl-Lysine, g	0.41	0.46	0.5	0.54	0.59
DCP, g	2.1	2.1	2.1	2.1	2.1
Shell, g	0.3	0.3	0.3	0.3	0.3
Premix, g	0.25	0.25	0.25	0.25	0.25
Salt, g	0.48	0.48	0.48	0.48	0.48
Nutrient value					
ME, kcal/kg	2,930	2,930	2,929	2,928	2,926
CP, %	18	18	18	18	18
Lysine, %	0.9	0.95	0.99	1.03	1.08
Ca, %	1.2	1.2	1.2	1.2	1.2
P, %	0.78	0.78	0.78	0.78	0.78

Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding 5%, 10%, 15%, and 20% lysine compared with control treatment; DCP: Dicarboxate phosphorus; CP: crude protein; ca: Calcium; P: phosphorus.

B. Feed intake, body weight and carcass composition

Feed intake was recorded daily by weighing feed supply and refusal. Body weight was taken every week at 6 a.m. before chickens were fed. Feed conversion ratio was calculated as feed intake per unit of body weight. For carcass composition, 12 chickens (per treatment) were slaughtered at 8 weeks of age to determine the carcass, breast meat and leg meat ratios. Breast meat nutritional values (dry matter, crude protein, fat and ash content) were analyzed by Weende's proximate method [2] and amino acid composition was determined using the high-pressure liquid chromatography. Internal organs such as spleen, thymus and bursa of Fabricius were separated and weighed and the relative organ weight was calculated as percentage of body weight.

C. Antibody response analysis

A total of 6 blood samples in each treatment were collected from 6 birds via their hearts to determine maternal antibody levels against the bird flu disease. Determining antibody against the bird flu disease was done using haemagglutination method (HA) titration as suggested by [25].

D. Statistical analysis

Data on growth performance, bodyweight, carcass composition and immunological responses were analyzed by one-way of variance (ANOVA) in [12]. The differences between experimental treatments were examined with Tukey's method.

III. RESULTS AND DISCUSSION

A. Growth performance and feed utilization

Body weight gain (BWG) and feed conversion ratio (FCR) of birds were significantly influenced ($P < 0.05$) by the different diets (Table 4), of which BWG was highest in the Ctrl+10 (52.4 g/bird/week) but was lowest in Ctrl (43.62 g/bird/week) diet. Similarly, the feed conversion of birds was most efficient in Ctrl+10 (2.09 kg feed/kg weight gain), compared with other treatments. However, there was no significant difference regarding to feed intake of chickens among treatments. The present results partly supported the observations of [11] and [6] that lysine supplementation above NRC requirement [16] in broiler diets improved final body weight, total weight gain and feed intake.

Previously, the finding of [14] also identified a positive linearity of the relationship between lysine requirement and dietary protein level, which supported the view that lysine requirement for maximum productivity (maximum weight gain and feed efficiency) was in the range of 1.26 to 1.33% and this was above the NRC requirement of 1%. Moreover, [5] found that lysine supplementation significantly improved the live body weight and feed conversion efficiency. Supportably, [27] pointed out that the poor live weight performance of broilers fed with basal diets and the responses to supplemental L-lysine indicated that the basal diets were severely deficient in lysine. From the present work, it was also indicated that supplementary levels more than NRC requirement (>5%) was appropriate for enhancing growth rate of Ac chicken. This could be explained by the balance of amino acid content in the diets, where lysine is the basic unit to synthesize protein, it affects directly protein synthesis in the body of birds, and thereby improved growth rate of birds.

TABLE 4
Effects of dietary lysine on birds' body weight and feed conversion ratio

Variables	Treatments					P
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20	
Initial BW, g/bird	21.3	22.0	21.0	22.6	21.6	n.s
FI, g/bird/week	99.17	104.6	111.9	104.5	103.9	0.221
BWG, g/bird/week	43.62 ^b	48.0 ^{ab}	52.4 ^a	48.8 ^{ab}	48.1 ^{ab}	0.021
FCR	2.29 ^a	2.16 ^b	2.09 ^b	2.12 ^b	2.13 ^b	0.000

FCR: feed conversion ratio; BW: bodyweight; BWG: bodyweight gain; FI: feed intake; Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding 5%, 10%, 15%, and 20% lysine compared with control treatment; ns: non-significant;

^{a,b} means with different superscripts within a row are significantly different at $P < 0.05$.

B. Carcass composition and nutritional value of breast muscle

TABLE 5
Effects of dietary lysine on birds' carcass performance

Parameters	Treatments					P
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20	
Body weight, g	395	400	397	399	404	0.941
Carcass, g	276	280	280	280	281	0.940
Carcass ratio, %	69.9	70.1	70.4	70.2	69.6	0.753
Breast muscle ratio, %	19.6	19.7	18.75	19.53	19.9	0.144
Thigh muscle ratio, %	20.5	20.3	20.4	20.9	20.1	0.511

Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding 5%, 10%, 15%, and 20% lysine compared with control treatment.

The composition of carcass and nutritional value of breast muscle of Ac chicken was not different among diets ($P > 0.05$) (Table 5 and Table 6). The present output agreed with the finding of [23], which stated that carcass weight and breast muscle yield were not affected by dietary lysine. [20] also reported that lysine had no effects on breast and leg muscle weight. In contrast, [13] and [1] presented that breast meat yield reduced significantly from 1.05% to 0.85% when lysine content was limited. Moreover, [24] and [19] showed that excessive lysine increased muscle weights and breast muscle weight. In term of nutritional value, dry matter and protein content of breast meat were similar to the report of [26]. The results were consistent with findings of [7], who concluded that crude protein and crude fat of thigh meat were not influenced by the replacement of duckweed for soybean and rice bran in chicken diet. Similar results were also found in amino acid composition, where no effect of lysine was observed.

TABLE 6
Effects of lysine on nutritional value and amino acid composition of breast muscle

Parameters	Treatments					P
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20	
Nutritional value						
Dry matter (%)	25.75±1.13	25.76±0.42	26.03±0.66	26.06±1.11	25.22±0.89	0.488
Crude protein (%)	23.68±2.21	23.20±1.19	23.52±0.51	23.37±1.11	22.32±0.82	0.430
Fat (%)	3.26±0.92	3.12±0.57	2.54±0.76	2.37±1.04	2.85±0.92	0.359
Ash (%)	2.01±1.13	1.32±0.13	1.65±0.74	1.33±0.06	1.25±0.21	0.206
Amino acid (g/kg)						
Aspartic	17.73±1.37	17.12±1.34	16.86±0.8	17.35±1.47	18.55±0.51	0.146
Serine	17.48±0.71	18.84±5.17	16.28±0.68	17.02±1.71	17.3±1.12	0.525
Glutamic	32.08±3.18	32.42±2.43	31.12±1.20	31.82±3.37	32.62±1.95	0.862
Glycine	28.96±5.82	28.16±7.33	26.34±4.41	25.35±4.65	22.85±1.57	0.286
Histidine	23.88±2.39	26.56±2.61	25.1±2.82	26.67±3.01	25.63±1.58	0.320
Threonine	18.54±0.54	18.71±2.21	18.71±1.59	16.05±4.35	17.76±1.26	0.275
Proline	15.22±1.76	14.34±1.15	14.57±1.48	14.7±1.45	14.05±1.01	0.675
Cystine	9.10±1.60	7.08±2.04	9.27±1.94	9.97±1.34	9.45±1.55	0.067
Tyrosine	27.22±3.03	27.93±2.01	25.76±2.54	28.21±4.43	26.83±5.87	0.818
Valine	20.12±1.81	19.63±0.65	19.62±0.74	20.1±1.65	20.6±1.33	0.694
Methionine	14.16±2.09	11.47±4.76	14.03±2.04	15.47±1.56	14.33±1.67	0.165
Lysine	13.95±3.14	12.4±0.79	14.77±1.3	12.42±0.84	13.21±2.16	0.163
Leucine	29.01±2.63	29.92±0.98	29.43±0.77	30.4±3.3	30.77±1.94	0.630
Phenylalanine	30.57±3.87	32.64±2.11	29.95±2.78	32.88±4.7	30.99±1.82	0.448
Tryptophane	2.13±0.13	2.05±0.04	2.14±0.09	2.05±0.15	1.98±0.08	0.072

Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding 5%, 10%, 15%, and 20% lysine compared with control treatment.

C. Immunological response to influenza vaccination

Evidences in the report of [8] and [9] showed that a dietary deficiency of lysine limited the synthesis of proteins and the proliferation of lymphocytes, as well as impaired immunological responses in birds, resulting in increases in morbidity and mortality in response to infection. There were also findings stating that an inadequate intake of dietary lysine reduced antibody responses and cell-mediated immunity in chickens [4]. In addition, lysine and methionine supplementation to the diet has improved the immunity of chickens against potential diseases [21]. In the present study, there was no significant difference on birds' spleen ratio between diets; however, Fabricius and thymus ratio of birds was statistically influenced ($P < 0.05$) by different amounts of dietary lysine (Table 7). In the Ctrl+20, Ctrl+15, and Ctrl+10, the ratios of Fabricius (0.146%, 0.141%, and 0.134%, respectively) and thymus (0.522%, 0.542%, and 0.536%, respectively) was higher than that of Ctrl with 0.104% of Fabricius and 0.382% of thymus.

As dietary lysine contents increased from 1% to 1.15% and from 0.9 to 1.03 in stages 0-4 and 5-8 weeks of age, respectively, the ratio of Fabricius and thymus gradually increased. Lysine is an essential amino acid that is necessary to produce proteins including antibodies; thus, adequate dietary levels of lysine is needed to support optimum efficacy of the immune system. Although the value of \log_2 HI titre of the Ctrl treatment tended to be lower than those of the other treatments, the antibody titre against avian influenza at 28 and 56 day-old were not significantly different among treatments. [11] reported that increased lysine content in diets of broilers in excess of NRC recommendation could improve immune system functions against Newcastle disease. Therefore, an appropriate vaccination regime should be taken into consideration for Ac chicken to fully achieve the efficacy of immune function.

TABLE 7
Effects of lysine on birds' immune responses

Variables	Treatments					P
	Ctrl	Ctrl+5	Ctrl+10	Ctrl+15	Ctrl+20	
Spleen ratio, %	0.09±0.013	0.10±0.023	0.105±0.023	0.104±0.02	0.103±0.019	0.815
Fabricius ratio, %	0.104 ^b ±0.02	0.123 ^{ab} ±0.01	0.134 ^a ±0.01	0.141 ^a ±0.02	0.146 ^a ±0.02	0.000
Thymus ratio, %	0.382 ^b ±0.07	0.475 ^{ab} ±0.03	0.536 ^a ±0.08	0.542 ^a ±0.06	0.522 ^a ±0.06	0.001
H ₅ N ₁ antibody, 28d*	6.81±1.05	7.5±0.75	7.64±0.63	7.08±1.38	7.28±0.85	0.838
H ₅ N ₁ antibody, 56d*	6.75±0.90	7.72±0.25	7.5±0.43	7.33±0.76	7.18±0.71	0.482

Ctrl: control treatment; Ctrl+5, Ctrl+10, Ctrl+15, and Ctrl+20: treatments adding 5%, 10%, 15%, and 20% lysine compared with control treatment; *: log₂ HI titre; ^{a,b} means with different superscripts within a row are significantly different at P < 0.05.

IV. CONCLUSIONS

The present study demonstrated that dietary lysine supplementation at 10% improved growth performance and feed conversion ratio but carcass composition were not influenced. The ratios of spleen and thymus were improved but immunological responses to influenza vaccination remained unchanged despite the increased lysine content in the diet.

REFERENCES

- [1] Acar N., Moran E. T., and Bilgili S. F. 1991. Live performance and carcass yield of male broilers from two commercial strain crosses receiving rations containing lysine below and above the established requirement between six and eight weeks of age. *Poultry Science* 70: 2315-2321.
- [2] AOAC 1990. Official methods of analysis, 15th Edition. Association of official analytical chemists. Washington, DC. Volume 1: 69-90.
- [3] Baker D. H., Batal A. B., Parr T. M., Augspurger N. R., and Parsons C. M. 2002. Ideal ratio (relative to lysine) of tryptophan, threonine, isoleucine and valine for chicks during the second and third posthatch. *Poultry Science* 8: 485-494.
- [4] Chen C., Sander J. E., and Dale N. M. 2003. The effect of dietary lysine deficiency on the immune response to Newcastle disease vaccination in chickens. *Avian Dis* 47: 1346-1351.
- [5] Corzo A., Kidd M. T., Burnham D. J., Miller E. R., Branton S. L., and Gonzalez-Esquerria R. 2005. Dietary amino acid density effects on growth and carcass of broilers differing in strain cross and sex. *Journal Applied Poultry Research* 14: 1-9.
- [6] Faluyi O. B., Agbede J. O., and Adebayo I. A. 2015. Growth performance and immunological response to Newcastle disease vaccination of broiler chickens fed lysine supplement diets. *Journal Veterinary Medicine Animal Health* 7: 77-84.
- [7] Khang N. T. K. and Ogle B. 2004. Effects of replacing roasted soya beans by broken rice and duckweed on performance of growing Tau Vang chickens confined on-station and scavenging on-farm. *Livestock Research Rural Development* 16 (8).
- [8] Kidd M. T., Kerr B. J., and Anthony N. B. 1997. Dietary interactions between lysine and threonine in broilers. *Poultry Science* 76: 608-614.
- [9] Konashi S., Takahashi K. and Akiba Y. 2000. Effects of dietary essential amino acid deficiencies on immunological variables in broiler chickens. *British Journal Nutrition* 83: 449-456.
- [10] Labadan M. C., Hsu K. N. and Austic R. E. 2001. Lysine and arginine requirements of broiler chickens at two-to-three-week intervals to eight weeks of age. *Poultry Science* 80: 599-606.
- [11] Mehrdad B. 2012. Effect of excess lysine and methionine on immune system and performance of broilers. *Annals Biology Science* 3: 3218-3224.
- [12] Minitab, 2000. Minitab Reference Manual. PC Version, Release 13.2. Minitab Inc., State College, PA.
- [13] Moran E. T. and Bilgili S. F. 1990. Processing losses, carcass quality, and meat yields of broiler chickens receiving diets marginally deficient to adequate in lysine prior to marketing. *Poultry Science* 69: 702-710.
- [14] Morris T. R. and Abebe S. 1990. Note on the effect of protein concentration on response to dietary lysine by chicks. *British Poultry Science* 31: 267-272.
- [15] Murray R. K., Granner D. K., Mayes P. A., and Rodwell V. W. 1998. Harper's illustrated biochemistry. Appleton Lana; Norwalk, CT, USA. pp. 610-617.
- [16] National Research Council, 1994. Nutrient Requirements of Poultry, 9th edition. National Academy Press, Washington DC.
- [17] Phuoc T. V., Dung N. N. X., and Manh L. H. 2015. Effect of dietary protein and every levels on performance and egg quality of Ac chicken at the first laying stage. *National Proceedings on Animal and Veterinary Sciences*. Can Tho University, Vietnam, pp.335-341 (in Vietnamese).
- [18] Phuong T. T. M. and Thien N. V. 2008. Ac chicken production in Vietnam. In Proceedings of the 7th RBI global conference on the conservation of animal genetic resources "impact of globalization on the animal genetic resources", Ha Noi, Vietnam, pp. 140-145.
- [19] Rezaei M., Moghaddam H. N., Reza J. P., and Kermanshahi H. 2004. The effects of dietary protein and lysine levels on broiler performance, carcass characteristics and N excretion. *Journal Poultry Science* 3: 148-152.
- [20] Rostagno H. S. and Pack M. 1995. Growth and breast meat responses of different broiler strains to dietary lysine. In Proceedings of the 10th European Symposium on Poultry Nutrition, Antalya, Turkey, pp. 260-262.
- [21] Rubin L. L., Ribeiro A. L. M., Canal C. W., Silva I. C., Trevizan L., Vogt L. K., Pereira R. A., and Lacerda L. 2007. Influence of sulfur amino acid levels in diets of broiler chickens submitted to immune stress. *Brazilian Journal Poultry Science* 9: 305-311.
- [22] Sterling K. G., Pest G., and Bakall M. R. I. 2008. Performance of different broiler genotypes fed diets with varying levels of dietary crude protein and lysine. *Poultry Science* 85: 1045-1054.

International Journal of Emerging Technology and Advanced Engineering

Website: www.ijetae.com (ISSN 2250-2459, ISO 9001:2008 Certified Journal, Volume 7, Issue 7, July 2017)

- [23] Tang M. Y, Ma Q. G., Chen X. D., and Ji C. 2007. Effects of dietary metabolizable energy and lysine on carcass characteristics and meat quality in Arbor Acres broilers. *Asian-Australasian Journal Animal Science* 20: 1865-1873.
- [24] Tesseraud S., Temin S., Le Bihan-Duval E., and Chagneau A. M. 2001. Increased responsiveness to dietary lysine deficiency of pectoralis major muscle protein turnover in broilers selected on breast development. *Journal Animal Science* 79: 927-933.
- [25] Thayer S. G. and Beard C. W. 1998. Serological procedure in: A laboratory manual for the isolation and identification of avian pathogens, 4th Ed. American Association of Avian Pathologist, Pennsylvania, USA, pp. 255-258.
- [26] Thieu P. C., Tieu H. V., Su V. V., Son H. L., Son V. N., and Nhan T. K. 2008. Conservation and development gene source of H'mong chicken. In Proceedings of the 7th RBI global conference on the conservation of animal genetic resources "impact of globalization on the animal genetic resources", Ha Noi, Vietnam, pp. 79-83.
- [27] Usama A., Muhammad A., Abdul S. M., and Zhirong J. 2007. Lysine requirement of broiler chickens fed low-density diets under tropical conditions. *Asian-Australasian Journal Animal Science* 20: 939-943.